

# STAT 401A - Statistical Methods for Research Workers

## Modeling assumptions

Jarad Niemi (Dr. J)

Iowa State University

last updated: September 15, 2014

# Normality assumptions


In the paired t-test, we assume

$$D_i \stackrel{iid}{\sim} N(\mu, \sigma^2).$$


In the two-sample t-test, we assume

$$Y_{ij} \stackrel{ind}{\sim} N(\mu_j, \sigma^2).$$

**Paired t-test**



**Two-sample t-test**



## Normality assumptions

In the paired t-test, we assume

$$D_i \stackrel{iid}{\sim} N(\mu, \sigma^2).$$


In the two-sample t-test, we assume

$$Y_{ij} \stackrel{ind}{\sim} N(\mu_j, \sigma^2).$$


Key features of the normal distribution assumption:

- Centered at the mean (expectation)  $\mu$
- Standard deviation describes the spread
- Symmetric around  $\mu$  (no skewness)
- Non-heavy tails, i.e. outliers are rare (no kurtosis)


# Normality assumptions




# Kurtosis (heavy-tailedness)




# Kurtosis (heavy-tailedness)



# Kurtosis (heavy-tailedness)




# Kurtosis (heavy-tailedness)






# Skewness



# Samples from skewed distributions



# Robustness

## Definition

A statistical procedure is **robust to departures from a particular assumption** if it is valid even when the assumption is not met.

**Remark** If a 95% confidence interval is robust to departures from a particular assumption, the confidence interval should cover the true value about 95% of the time.


## Robustness to skewness and kurtosis

Percentage of 95% confidence intervals that cover the true difference in means in an equal-sample two-sample t-test with non-normal populations (where the distributions are the same other than their means).


| sample size | strongly skewed | moderately skewed | mildly skewed | heavy-tailed | short-tailed |
|-------------|-----------------|-------------------|---------------|--------------|--------------|
| 5           | 95.5            | 95.4              | 95.2          | 98.3         | 94.5         |
| 10          | 95.5            | 95.4              | 95.2          | 98.3         | 94.6         |
| 25          | 95.3            | 95.3              | 95.1          | 98.2         | 94.9         |
| 50          | 95.1            | 95.3              | 95.1          | 98.1         | 95.2         |
| 100         | 94.8            | 95.3              | 95.0          | 98.0         | 95.6         |

# Differences in variances

Normal distribution



# Differences in variances



## Robustness to differences in variances

Percentage of 95% confidence intervals that cover the true difference in means in an equal-sample two-sample t-test ( $r = \sigma_1/\sigma_2$ ).

| n1  | n2  | r=1/4 | r=1/2 | r=1  | r=2  | r=4  |
|-----|-----|-------|-------|------|------|------|
| 10  | 10  | 95.2  | 94.2  | 94.7 | 95.2 | 94.5 |
| 10  | 20  | 83.0  | 89.3  | 94.4 | 98.7 | 99.1 |
| 10  | 40  | 71.0  | 82.6  | 95.2 | 99.5 | 99.9 |
| 100 | 100 | 94.8  | 96.2  | 95.4 | 95.3 | 95.1 |
| 100 | 200 | 86.5  | 88.3  | 94.8 | 98.8 | 99.4 |
| 100 | 400 | 71.6  | 81.5  | 95.0 | 99.5 | 99.9 |

# Outliers

## Definition

A statistical procedure is **resistant** if it does not change very much when a small part of the data changes, perhaps drastically.

Identify outliers:

- 1 If recording errors, fix.
- 2 If outlier comes from a different population, remove and report.
- 3 If results are the same with and without outliers, report with outliers.
- 4 If results are different, use resistant analysis or report both analyses.



# Common ways for independence to be violated

- Cluster effect
  - e.g. pigs in a pen
- Serial effect
  - e.g. measurements in time with drifting scale
- Spatial effect
  - e.g. corn yield plots (drainage)

# Common transformations for data

From: [http://en.wikipedia.org/wiki/Data\\_transformation\\_\(statistics\)](http://en.wikipedia.org/wiki/Data_transformation_(statistics))

## Definition

In statistics, **data transformation** refers to the application of a deterministic mathematical function to each point in a data set that is, each data point  $y_i$  is replaced with the transformed value  $z_i = f(y_i)$ , where  $f$  is a function.

The most common transformations are

- If  $y$  is a proportion, then  $f(y) = \sin^{-1}(\sqrt{y})$ .
- If  $y$  is a count, then  $f(y) = \sqrt{y}$ .
- If  $y$  is positive and right-skewed, then  $f(y) = \log(y)$ , the *natural logarithm* of  $y$ .

**Remark** Since  $\log(0) = -\infty$ , the logarithm cannot be used directly when some  $y_i$  are zero. In these cases, use  $\log(y + c)$  where  $c$  is something small relative to your data, e.g. half of the minimum non-zero value.

# Log transformation

Consider two-sample data and let  $z_{ij} = \log(y_{ij})$ . Now, run a two-sample t-test on the  $z$ 's. Then we assume

$$Z_{ij} \stackrel{ind}{\sim} N(\mu_j, \sigma^2)$$

and the quantity  $\bar{Z}_2 - \bar{Z}_1$  estimates the “difference in population means on the (natural) log scale”. The quantity  $\exp(\bar{Z}_2 - \bar{Z}_1) = e^{\bar{Z}_2 - \bar{Z}_1}$  estimates

$$\frac{\text{Median of population 2}}{\text{Median of population 1}}$$

on the original scale or, equivalently, it estimates the **multiplicative effect** of moving from population 1 to population 2.

## Log transformation interpretation

If we have a randomized experiment:

**Remark** It is estimated that the response of an experimental unit to treatment 2 will be  $\exp(\bar{Z}_2 - \bar{Z}_1)$  times as large as its response to treatment 1.

If we have an observational study:

**Remark** It is estimated that the median for population 2 is  $\exp(\bar{Z}_2 - \bar{Z}_1)$  times as large as the median for population 1.

## Confidence intervals with log transformation

If  $z_{ij} = \log(y_{ij})$  and we assume

$$Z_{ij} \stackrel{\text{ind}}{\sim} N(\mu_j, \sigma^2),$$

then a  $100(1 - \alpha)\%$  two-sided confidence interval for  $\mu_2 - \mu_1$  is

$$(L, U) = \bar{Z}_2 - \bar{Z}_1 \pm t_{n_1+n_2-2}(1 - \alpha/2)SE(\bar{Z}_2 - \bar{Z}_1).$$


A  $100(1 - \alpha)\%$  confidence interval for

$$\frac{\text{Median of population 2}}{\text{Median of population 1}}$$


is  $(e^L, e^U)$ .

# Miles per gallon data

Untransformed:




Logged:




# Miles per gallon data

Untransformed:



Logged:



## Equal variances?

We might also be concerned about the assumption of equal variances.

Untransformed:

| country | n   | mean  | sd   |
|---------|-----|-------|------|
| Japan   | 79  | 30.48 | 6.11 |
| US      | 249 | 20.14 | 6.41 |

the ratio of sample standard deviations is around 1.05 and there are 3 times as many observations in the US.

Logged:

| country | n   | mean | sd   |
|---------|-----|------|------|
| Japan   | 79  | 3.40 | 0.21 |
| US      | 249 | 2.96 | 0.31 |

Now the ratio of standard deviations is 1.5 which argues for not using the logarithm.



## 95% two-sample CI for the ratio by hand

| country | n   | mean | sd   |
|---------|-----|------|------|
| Japan   | 79  | 3.40 | 0.21 |
| US      | 249 | 2.96 | 0.31 |

Choose group 2 to be Japan and group 1 to be the US:

$$\begin{aligned}
 \alpha &= 0.05 \\
 n_1 + n_2 - 2 &= 249 + 79 - 2 = 326 \\
 t_{n_1+n_2-2}(1 - \alpha/2) &= t_{326}(0.975) = 1.96 \\
 \bar{Z}_2 - \bar{Z}_1 &= 3.40 - 2.96 = 0.44 \\
 s_p &= \sqrt{\frac{(n_1-1)s_1^2 + (n_2-1)s_2^2}{n_1+n_2-2}} = \sqrt{\frac{(249-1)0.31^2 + (79-1)0.21^2}{249+79-2}} = 0.29 \\
 SE(\bar{Z}_2 - \bar{Z}_1) &= s_p \sqrt{\frac{1}{n_1} + \frac{1}{n_2}} = 0.29 \sqrt{\frac{1}{249} + \frac{1}{79}} = 0.037
 \end{aligned}$$

Thus a 95% two-sided confidence interval for the difference (on the log scale) is

$$\begin{aligned}
 (L, U) &= \bar{Z}_2 - \bar{Z}_1 \pm t_{n_1+n_2-2}(1 - \alpha/2)SE(\bar{Z}_2 - \bar{Z}_1) \\
 &= 0.44 \pm 1.96 \times 0.037 \\
 &= (0.37, 0.51)
 \end{aligned}$$

and a 95% two-sided confidence interval for the ratio (on the original scale) is

$$(e^L, e^U) = (e^{0.37}, e^{0.51}) = (1.45, 1, 67)$$

# Using R for t-test using logarithms

```
t = t.test(log(mpg)~country, d, var.equal=TRUE)
t$estimate # On log scale

mean in group Japan    mean in group US
      3.396              2.955

exp(t$estimate) # On original scale

mean in group Japan    mean in group US
      29.85              19.21

exp(t$estimate[1]-t$estimate[2]) # Ratio of medians (Japan/US)

mean in group Japan
      1.554

exp(t$conf.int) # Confidence interval for ratio of medians

[1] 1.445 1.672
attr(,"conf.level")
[1] 0.95
```

## SAS code for t-test using logarithms

```
DATA mpg;
  INFILE 'mpg.csv' DELIMITER=', ' FIRSTOBS=2;
  INPUT mpg country $;

PROC TTEST DATA=mpg TEST=ratio;
CLASS country;
VAR mpg;
run;
```

# SAS output for t-test using logarithms

## The TTEST Procedure

Variable: mpg

| country     | N   | Geometric Mean | Coefficient of Variation | Minimum | Maximum |
|-------------|-----|----------------|--------------------------|---------|---------|
| Japan       | 79  | 29.8525        | 0.2111                   | 18.0000 | 47.0000 |
| US          | 249 | 19.2051        | 0.3147                   | 9.0000  | 39.0000 |
| Ratio (1/2) |     | 1.5544         | 0.2928                   |         |         |

| country     | Method        | Geometric Mean | 95% CL Mean |         | Coefficient of Variation | 95% CL CV |        |
|-------------|---------------|----------------|-------------|---------|--------------------------|-----------|--------|
| Japan       |               | 29.8525        | 28.4887     | 31.2817 | 0.2111                   | 0.1820    | 0.2514 |
| US          |               | 19.2051        | 18.4825     | 19.9560 | 0.3147                   | 0.2882    | 0.3467 |
| Ratio (1/2) | Pooled        | 1.5544         | 1.4452      | 1.6719  | 0.2928                   | 0.2712    | 0.3183 |
| Ratio (1/2) | Satterthwaite | 1.5544         | 1.4636      | 1.6508  |                          |           |        |

| Method        | Coefficients |        |         |         |
|---------------|--------------|--------|---------|---------|
|               | of Variation | DF     | t Value | Pr >  t |
| Pooled        | Equal        | 326    | 11.91   | <.0001  |
| Satterthwaite | Unequal      | 193.33 | 14.46   | <.0001  |

## Equality of Variances

| Method   | Num DF | Den DF | F Value | Pr > F |
|----------|--------|--------|---------|--------|
| Folded F | 248    | 78     | 2.17    | 0.0001 |

## Conclusion

Japanese median miles per gallon is 1.55 [95% CI (1.46,1.65)] times as large as US median miles per gallon.

OR

Japanese median miles per gallon is 55% [95% CI (46%,65%)] larger than US median miles per gallon.

## Unequal standard deviations

The two-sample t-test tools assume either

$$Y_{ij} \stackrel{ind}{\sim} N(\mu_j, \sigma^2) \quad \text{or} \quad Z_{ij} \stackrel{ind}{\sim} N(\mu_j, \sigma_j^2)$$

depending on whether we were working on the original scale ( $Y$ ) or log scale ( $Z$ ), respectively.

But what if we don't believe the variances in the two populations are equal, e.g. in the log transformed miles per gallon data set?


Instead compare

$$Y_{ij} \stackrel{ind}{\sim} N(\mu_j, \sigma_j^2) \quad \text{or} \quad Z_{ij} \stackrel{ind}{\sim} N(\mu_j, \sigma_j^2),$$

i.e. the populations have unequal variances. But still test  $H_0 : \mu_1 = \mu_2$  vs  $H_1 : \mu_1 \neq \mu_2$  or construct a confidence interval for  $\mu_2 - \mu_1$ .

# Visualization of two normals with unequal standard deviations

```
curve(dnorm, -3, 6, lwd=2)  
curve(dnorm(x, 2, 2), lwd=2, col=2, lty=2, add=TRUE)
```



# Welch's SE with Satterthwaite's approximation to df

Estimate of  $(\mu_2 - \mu_1)$ :

$$\bar{Y}_2 - \bar{Y}_1$$

Standard error:

$$SE_W(\bar{Y}_2 - \bar{Y}_1) = \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}$$

Degrees of freedom using the Satterthwaite's approximation:

$$df_W = \frac{SE_W(\bar{Y}_2 - \bar{Y}_1)^4}{\frac{SE(\bar{Y}_2)^4}{n_2 - 1} + \frac{SE(\bar{Y}_1)^4}{n_1 - 1}}$$

where

$$SE(\bar{Y}_2) = \frac{s_2}{\sqrt{n_2}} \quad \text{and} \quad SE(\bar{Y}_1) = \frac{s_1}{\sqrt{n_1}}$$

(which is the same formula as in the paired t-test)



## Welch's t-test and CI

Welch's t-test has test statistic:

$$t = \frac{\text{(Estimate-Parameter)}}{\text{SE(Estimate)}} = \frac{\bar{Y}_2 - \bar{Y}_1 - (\mu_2 - \mu_1)}{SE_W(\bar{Y}_2 - \bar{Y}_1)}$$

which has a  $t$  distribution with (approximately)  $df_W$  degrees of freedom if the null hypothesis is true. Calculate the pvalue

- Two-sided ( $H_1 : \mu_2 \neq \mu_1$ ):  $p = 2P(t_{df_W} < -|t|)$
- One-sided ( $H_1 : \mu_2 > \mu_1$ ):  $p = P(t_{df_W} < -t)$
- One-sided ( $H_1 : \mu_2 < \mu_1$ ):  $p = P(t_{df_W} < t)$

Two-sided  $100(1 - \alpha)\%$  confidence interval for  $\mu_2 - \mu_1$ :

$$\bar{Y}_2 - \bar{Y}_1 \pm t_{df_W}(1 - \alpha/2)SE_W(\bar{Y}_2 - \bar{Y}_1)$$

## Are the variances equal?

Suppose

$$Y_{ij} \stackrel{\text{ind}}{\sim} N(\mu_j, \sigma_j^2)$$

and you want to test  $H_0 : \sigma_1 = \sigma_2$  vs  $H_1 : \sigma_1 \neq \sigma_2$ .

You can use an  $F$ -test and its associated pvalue. If the pvalue is small, e.g. less than 0.05, then we reject  $H_0$ . If the pvalue is not small, then we fail to reject  $H_0$ , but this does not mean the variances are not equal.

(Section 4.5.3) discusses another approach called Levene's test

# Welch's test and CI using R

```
var.test(mpg~country,d) # F-test
```

F test to compare two variances

```
data: mpg by country
F = 0.9066, num df = 78, denom df = 248, p-value = 0.6194
alternative hypothesis: true ratio of variances is not equal to 1
95 percent confidence interval:
 0.6423 1.3246
sample estimates:
ratio of variances
      0.9066
```

```
(t=t.test(mpg~country, d, var.equal=FALSE))
```

Welch Two Sample t-test

```
data: mpg by country
t = 12.95, df = 136.9, p-value < 2.2e-16
alternative hypothesis: true difference in means is not equal to 0
95 percent confidence interval:
 8.758 11.915
sample estimates:
mean in group Japan    mean in group US
      30.48             20.14
```

## SAS code for two-sample t-test

```
DATA mpg;
  INFILE 'mpg.csv' DELIMITER=', ' FIRSTOBS=2;
  INPUT mpg country $;

PROC TTEST DATA=mpg;
  CLASS country;
  VAR mpg;
RUN;
```

# SAS output for t-test

## The TTEST Procedure

Variable: mpg

| country    | N   | Mean    | Std Dev | Std Err | Minimum | Maximum |
|------------|-----|---------|---------|---------|---------|---------|
| Japan      | 79  | 30.4810 | 6.1077  | 0.6872  | 18.0000 | 47.0000 |
| US         | 249 | 20.1446 | 6.4147  | 0.4065  | 9.0000  | 39.0000 |
| Diff (1-2) |     | 10.3364 | 6.3426  | 0.8190  |         |         |

| country    | Method        | Mean    | 95% CL Mean |         | Std Dev | 95% CL Std Dev |        |
|------------|---------------|---------|-------------|---------|---------|----------------|--------|
| Japan      |               | 30.4810 | 29.1130     | 31.8491 | 6.1077  | 5.2814         | 7.2429 |
| US         |               | 20.1446 | 19.3439     | 20.9452 | 6.4147  | 5.8964         | 7.0336 |
| Diff (1-2) | Pooled        | 10.3364 | 8.7252      | 11.9477 | 6.3426  | 5.8909         | 6.8699 |
| Diff (1-2) | Satterthwaite | 10.3364 | 8.7576      | 11.9152 |         |                |        |

| Method        | Variances | df     | t Value | Pr >  t |
|---------------|-----------|--------|---------|---------|
| Pooled        | Equal     | 326    | 12.62   | <.0001  |
| Satterthwaite | Unequal   | 136.87 | 12.95   | <.0001  |

## Equality of Variances

| Method   | Num df | Den df | F Value | Pr > F |
|----------|--------|--------|---------|--------|
| Folded F | 248    | 78     | 1.10    | 0.6194 |

```
var.test(log(mpg)~country,d)
```

F test to compare two variances

data: log(mpg) by country

F = 0.4617, num df = 78, denom df = 248, p-value = 0.0001055

alternative hypothesis: true ratio of variances is not equal to 1

95 percent confidence interval:

0.3271 0.6745

sample estimates:

ratio of variances

0.4617

```
(t = t.test(log(mpg)~country, d, var.equal=FALSE))
```

Welch Two Sample t-test

data: log(mpg) by country

t = 14.46, df = 193.3, p-value < 2.2e-16

alternative hypothesis: true difference in means is not equal to 0

95 percent confidence interval:

0.3809 0.5013

sample estimates:

| mean in group Japan | mean in group US |
|---------------------|------------------|
| 3.396               | 2.955            |

```
exp(t$conf.int)
```

```
[1] 1.464 1.651
```

```
attr( "conf.level" )
```

## SAS code for t-test using logarithms

```
DATA mpg;
  INFILE 'mpg.csv' DELIMITER=', ' FIRSTOBS=2;
  INPUT mpg country $;

PROC TTEST DATA=mpg TEST=ratio;
CLASS country;
VAR mpg;
run;
```

# SAS output for t-test using logarithms

## The TTEST Procedure

Variable: mpg

| country     | N   | Geometric Mean | Coefficient of Variation | Minimum | Maximum |
|-------------|-----|----------------|--------------------------|---------|---------|
| Japan       | 79  | 29.8525        | 0.2111                   | 18.0000 | 47.0000 |
| US          | 249 | 19.2051        | 0.3147                   | 9.0000  | 39.0000 |
| Ratio (1/2) |     | 1.5544         | 0.2928                   |         |         |

| country     | Method        | Geometric Mean | 95% CL Mean |         | Coefficient of Variation | 95% CL CV |        |
|-------------|---------------|----------------|-------------|---------|--------------------------|-----------|--------|
| Japan       |               | 29.8525        | 28.4887     | 31.2817 | 0.2111                   | 0.1820    | 0.2514 |
| US          |               | 19.2051        | 18.4825     | 19.9560 | 0.3147                   | 0.2882    | 0.3467 |
| Ratio (1/2) | Pooled        | 1.5544         | 1.4452      | 1.6719  | 0.2928                   | 0.2712    | 0.3183 |
| Ratio (1/2) | Satterthwaite | 1.5544         | 1.4636      | 1.6508  |                          |           |        |

| Method        | Coefficients |        |         |         |
|---------------|--------------|--------|---------|---------|
|               | of Variation | DF     | t Value | Pr >  t |
| Pooled        | Equal        | 326    | 11.91   | <.0001  |
| Satterthwaite | Unequal      | 193.33 | 14.46   | <.0001  |

## Equality of Variances

| Method   | Num DF | Den DF | F Value | Pr > F |
|----------|--------|--------|---------|--------|
| Folded F | 248    | 78     | 2.17    | 0.0001 |



# Summary

## Two-sample $t$ tools assumptions

- Normality
  - No skewness (take logs?)
  - No heavy tails
- Equal variances
  - Test: F-test or Levene's test
  - Use Welch's two-sample  $t$ -test and CI
- Independence (use random effects or avoid)
  - Cluster
  - Serial
  - Spatial