

R06a - Interpreting Regression p -values as Posterior Probabilities

STAT 587 (Engineering)
Iowa State University

March 30, 2021

Regression p -values

Recall the regression model

$$Y_i \stackrel{\text{ind}}{\sim} N(\mu_i, \sigma^2), \quad \mu_i = \beta_0 + \beta_1 X_{i,1} + \cdots + \beta_p X_{i,p}$$

A common hypothesis test is

$$H_0 : \beta_j = 0 \quad \text{versus} \quad H_A : \beta_j \neq 0$$

which has

$$p\text{-value} = 2P(T > |t|)$$

where $T \sim t_{n-(p+1)}$ and $t = \hat{\beta}_j / SE(\beta_j)$.

Example Regression Output

```
Call:
lm(formula = Speed ~ Conditions * log(NetToWinner), data = Sleuth3::ex0920)

Residuals:
    Min       1Q   Median       3Q      Max
-1.50551 -0.32127 -0.00219  0.35201  1.13026

Coefficients:
                Estimate Std. Error t value Pr(>|t|)
(Intercept)      33.23367    0.34584  96.095 < 2e-16 ***
ConditionsSlow    -2.04517    0.72404  -2.825  0.0056 **
log(NetToWinner)   0.27830    0.02942   9.458 5.88e-16 ***
ConditionsSlow:log(NetToWinner) 0.08664    0.06583   1.316  0.1908
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 0.4978 on 112 degrees of freedom
Multiple R-squared:  0.7015, Adjusted R-squared:  0.6935
F-statistic: 87.75 on 3 and 112 DF,  p-value: < 2.2e-16
```

Bayesian Posterior Probabilities

With prior $p(\beta, \sigma^2) \propto 1/\sigma^2$, we have

$$\beta_j | y \sim t_{n-(p+1)} \left(\hat{\beta}_j, SE(\beta_j)^2 \right).$$


Thus

$$P(\beta_j > 0 | y) = P \left(\frac{\beta_j - \hat{\beta}_j}{SE(\beta_j)} > \frac{0 - \hat{\beta}_j}{SE(\beta_j)} \middle| y \right) = P(T > -t)$$


which is very close to

$$p\text{-value} = 2P(T > |t|).$$


Visualizing Posterior Distribution



Visualizing Posterior Distribution



Visualizing Posterior Distribution



Interpreting Regression p -values as Posterior Probabilities

Suppose we have a p -value for $H_0 : \beta_j = 0$ vs $H_A : \beta_j \neq 0$. Then

- If $\hat{\beta}_j < 0$, then

$$P(\beta_j > 0|y) = p\text{-value}/2.$$

- If $\hat{\beta}_j > 0$, then

$$P(\beta_j < 0|y) = p\text{-value}/2.$$

Alternatively,

- If $\hat{\beta}_j < 0$, then

$$P(\beta_j < 0|y) = 1 - p\text{-value}/2.$$

- If $\hat{\beta}_j > 0$, then

$$P(\beta_j > 0|y) = 1 - p\text{-value}/2.$$

Example Interpretation

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	33.23	0.35	96.09	0.00
ConditionsSlow	-2.05	0.72	-2.82	0.01
log(NetToWinner)	0.28	0.03	9.46	0.00
ConditionsSlow:log(NetToWinner)	0.09	0.07	1.32	0.19

Intercept	$P(\beta_0 > 0 y) \approx 1$
ConditionsSlow	$P(\beta_1 < 0 y) \approx 0.99$
log(NetToWinner)	$P(\beta_2 > 0 y) \approx 1$
ConditionsSlow:log(NetToWinner)	$P(\beta_3 > 0 y) \approx 0.90$

Summary

Suppose we have a regression p -value for $H_0 : \beta_j = 0$ vs $H_A : \beta_j \neq 0$. Then

- If $\hat{\beta}_j < 0$, then

$$P(\beta_j < 0|y) = 1 - p\text{-value}/2.$$

- If $\hat{\beta}_j > 0$, then

$$P(\beta_j > 0|y) = 1 - p\text{-value}/2.$$