

R11 - ANOVA

HCI/PSYCH 522
Iowa State University

April 28, 2022

Outline

- One-way ANOVA
 - Mouse data
 - R code
 - Model
 - Mouse analysis
 - Bias in jury selection
- Two-way ANOVA
 - Seaweed grazer data
 - R code
 - Model
 - Seaweed grazer analysis
 - Pygmalion effect
- Summary
- Three-way ANOVA
 - Interactions

Mouse dataset

```
mouse <- read_csv('mouse.csv') %>%  
  mutate(Mouse = factor(Mouse),  
         Mouse = relevel(Mouse, ref="Dell"))  
head(mouse)
```


```
## # A tibble: 6 x 2  
##   Skill Mouse  
##   <dbl> <fct>  
## 1  35.5 Dell  
## 2  35.4 Dell  
## 3  34.9 Dell  
## 4  34.8 Dell  
## 5  33.8 Dell  
## 6  33.5 Dell
```

```
summary(mouse)
```

```
##      Skill      Mouse  
##  Min.   : 6.4   Dell      :49  
##  1st Qu.:31.8   Basilisk (Wired) :57  
##  Median :39.5   Mamba (Wired)    :71  
##  Mean   :38.8   Mamba (Wireless) :56  
##  3rd Qu.:46.9   Viper (Wired, light):60  
##  Max.   :54.6   Viper (Wired)    :56
```

Mouse graphically

```
ggplot(mouse, aes(x = Mouse, y = Skill)) + geom_jitter(width=0.1)
```



Regression model

```
m <- lm(Skill ~ Mouse, data = mouse)
summary(m)
```

```
##
## Call:
## lm(formula = Skill ~ Mouse, data = mouse)
##
## Residuals:
```

##	Min	1Q	Median	3Q	Max
##	-25.5167	-3.3857	0.8143	5.1833	10.0143

```
##
## Coefficients:
```

##		Estimate	Std. Error	t value	Pr(> t)
##	(Intercept)	27.402	0.954	28.722	< 2e-16 ***
##	MouseBasilisk (Wired)	5.289	1.301	4.065	5.95e-05 ***
##	MouseMamba (Wired)	14.895	1.240	12.009	< 2e-16 ***
##	MouseMamba (Wireless)	12.284	1.306	9.403	< 2e-16 ***
##	MouseViper (Wired, light)	17.715	1.286	13.776	< 2e-16 ***
##	MouseViper (Wired)	15.484	1.306	11.852	< 2e-16 ***

```
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 6.678 on 343 degrees of freedom
## Multiple R-squared:  0.4543, Adjusted R-squared:  0.4463
## F-statistic: 57.1 on 5 and 343 DF, p-value: < 2.2e-16
```

Confidence/credible intervals

```
coef(m)

##           (Intercept)      MouseBasilisk (Wired)      MouseMamba (Wired)      MouseMamba (Wireless)
##           27.402041           5.289187           14.895142           12.283673
## MouseViper (Wired, light)      MouseViper (Wired)
##           17.714626           15.483673

confint(m)

##           2.5 %      97.5 %
## (Intercept)      25.525547  29.278535
## MouseBasilisk (Wired)      2.730232  7.848142
## MouseMamba (Wired)      12.455599  17.334686
## MouseMamba (Wireless)      9.714178  14.853169
## MouseViper (Wired, light)  15.185417  20.243835
## MouseViper (Wired)      12.914178  18.053169
```

Regression model

```
em <- emmeans(m, pairwise ~ Mouse, adjust = "none")
confint(em)
```

```
## $emmeans
## Mouse          emmean    SE  df lower.CL upper.CL
## Dell           27.4 0.954 343    25.5    29.3
## Basilisk (Wired) 32.7 0.885 343    31.0    34.4
## Mamba (Wired)   42.3 0.793 343    40.7    43.9
## Mamba (Wireless) 39.7 0.892 343    37.9    41.4
## Viper (Wired, light) 45.1 0.862 343    43.4    46.8
## Viper (Wired)   42.9 0.892 343    41.1    44.6
##
## Confidence level used: 0.95
##
## $contrasts
## contrast          estimate    SE  df lower.CL upper.CL
## Dell - Basilisk (Wired)    -5.289 1.30 343    -7.848   -2.730
## Dell - Mamba (Wired)     -14.895 1.24 343   -17.335  -12.456
## Dell - Mamba (Wireless)  -12.284 1.31 343   -14.853   -9.714
## Dell - Viper (Wired, light) -17.715 1.29 343   -20.244  -15.185
## Dell - Viper (Wired)     -15.484 1.31 343   -18.053  -12.914
## Basilisk (Wired) - Mamba (Wired)    -9.606 1.19 343   -11.942   -7.270
## Basilisk (Wired) - Mamba (Wireless)  -6.994 1.26 343    -9.466   -4.523
## Basilisk (Wired) - Viper (Wired, light) -12.425 1.24 343   -14.855   -9.996
## Basilisk (Wired) - Viper (Wired)    -10.194 1.26 343   -12.666   -7.723
## Mamba (Wired) - Mamba (Wireless)     2.611 1.19 343     0.264    4.959
## Mamba (Wired) - Viper (Wired, light)  -2.819 1.17 343    -5.123   -0.516
## Mamba (Wired) - Viper (Wired)       -0.589 1.19 343    -2.936    1.759
## Mamba (Wireless) - Viper (Wired, light) -5.431 1.24 343    -7.872   -2.990
```

Regression model

```
summary(m)

##
## Call:
## lm(formula = Skill ~ Mouse, data = mouse)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -25.5167  -3.3857   0.8143   5.1833  10.0143
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)      27.402      0.954  28.722 < 2e-16 ***
## MouseBasilisk (Wired)    5.289      1.301   4.065 5.95e-05 ***
## MouseMamba (Wired)     14.895      1.240  12.009 < 2e-16 ***
## MouseMamba (Wireless)   12.284      1.306   9.403 < 2e-16 ***
## MouseViper (Wired, light) 17.715      1.286  13.776 < 2e-16 ***
## MouseViper (Wired)     15.484      1.306  11.852 < 2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 6.678 on 343 degrees of freedom
## Multiple R-squared:  0.4543, Adjusted R-squared:  0.4463
## F-statistic: 57.1 on 5 and 343 DF, p-value: < 2.2e-16
```


Analysis of variance (ANOVA)

```
anova(m)

## Analysis of Variance Table
##
## Response: Skill
##           Df Sum Sq Mean Sq F value    Pr(>F)
## Mouse      5  12734  2546.8   57.104 < 2.2e-16 ***
## Residuals 343  15297    44.6
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Analysis of variance (ANOVA)

```
anova(m)

## Analysis of Variance Table
##
## Response: Skill
##           Df Sum Sq Mean Sq F value    Pr(>F)
## Mouse      5  12734  2546.8   57.104 < 2.2e-16 ***
## Residuals 343  15297    44.6
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

ANOVA F-test: Comparison of models

Regression model with categorical variable:

$$Y_i \stackrel{ind}{\sim} N(\beta_0 + \beta_1 X_{i,1} + \cdots + \beta_p X_{i,p}, \sigma^2)$$

where $\beta_p, p > 0$ is the difference between mean response in the reference level compared to the level associated with the p th level

F-test:

- Reduced model: no categorical variable $\beta_1 = \cdots = \beta_p = 0$
- Full model: with categorical variable (see above)

ANOVA F-test: Summary

```
anova(m)

## Analysis of Variance Table
##
## Response: Skill
##           Df Sum Sq Mean Sq F value    Pr(>F)
## Mouse      5  12734  2546.8   57.104 < 2.2e-16 ***
## Residuals 343  15297    44.6
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

There is evidence of a difference in mean player skill using different mice ($F_{5,343} = 57, p \approx 0$).

YouTube videos

(hyperlinks)

Playlists:

- Probability
- Inference
- Regression
 - One-way ANOVA
 - F-tests

Bias in jury selection

```
case0502 <- Sleuth3::case0502 %>% mutate(Judge = relevel(Judge, ref="Spock's"))
head(case0502)
```


```
##   Percent  Judge
## 1     6.4 Spock's
## 2     8.7 Spock's
## 3    13.3 Spock's
## 4    13.6 Spock's
## 5    15.0 Spock's
## 6    15.2 Spock's
```

```
summary(case0502)
```

```
##      Percent      Judge
## Min.   : 6.40 Spock's:9
## 1st Qu.:19.95  A      :5
## Median :27.50  B      :6
## Mean   :26.58  C      :9
## 3rd Qu.:32.38  D      :2
## Max.   :48.90  E      :6
##                F      :9
```

Bias in jury selection - Plot

```
ggplot(case0502, aes(x = Judge, y = Percent)) + geom_jitter(width=0.1)
```



Bias in jury selection - One-way ANOVA

```
m <- lm(Percent ~ Judge, data = case0502)
anova(m)

## Analysis of Variance Table
##
## Response: Percent
##           Df Sum Sq Mean Sq F value    Pr(>F)
## Judge      6 1927.1   321.18   6.7184 6.096e-05 ***
## Residuals 39 1864.5    47.81
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Manuscript statement: There is evidence of a difference in mean percent women on juries amongst the judges ($F_{6,39} = 7, p \approx 0$).

Bias in jury selection - Model summary

```
summary(m)

##
## Call:
## lm(formula = Percent ~ Judge, data = case0502)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -17.320  -4.367  -0.250   3.319  14.780
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   14.622     2.305   6.344 1.72e-07 ***
## JudgeA        19.498     3.857   5.056 1.05e-05 ***
## JudgeB        18.994     3.644   5.212 6.39e-06 ***
## JudgeC        14.478     3.259   4.442 7.15e-05 ***
## JudgeD        12.378     5.405   2.290 0.027513 *
## JudgeE        12.344     3.644   3.388 0.001623 **
## JudgeF        12.178     3.259   3.736 0.000597 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 6.914 on 39 degrees of freedom
## Multiple R-squared:  0.5083, Adjusted R-squared:  0.4326
## F-statistic: 6.718 on 6 and 39 DF,  p-value: 6.096e-05
```

Bias in jury selection - Treatment vs Control

```
em <- emmeans(m, trt.vs.ctrl ~ Judge, adjust = "none")
confint(em)
```

```
## $emmeans
## Judge emmean SE df lower.CL upper.CL
## Spock's 14.6 2.30 39 9.96 19.3
## A 34.1 3.09 39 27.87 40.4
## B 33.6 2.82 39 27.91 39.3
## C 29.1 2.30 39 24.44 33.8
## D 27.0 4.89 39 17.11 36.9
## E 27.0 2.82 39 21.26 32.7
## F 26.8 2.30 39 22.14 31.5
##
## Confidence level used: 0.95
##
## $contrasts
## contrast estimate SE df lower.CL upper.CL
## A - Spock's 19.5 3.86 39 11.70 27.3
## B - Spock's 19.0 3.64 39 11.62 26.4
## C - Spock's 14.5 3.26 39 7.89 21.1
## D - Spock's 12.4 5.41 39 1.44 23.3
## E - Spock's 12.3 3.64 39 4.97 19.7
## F - Spock's 12.2 3.26 39 5.59 18.8
##
## Confidence level used: 0.95
```

Bias in jury selection - Custom contrast

Average of all other judge's percent women minus Spock's.

```
em <- emmeans(m, ~ Judge)
co <- contrast(em, list(`Mean(others) - Spock` = c(-7, rep(1,6))/7))
confint(co)

## contrast           estimate    SE df lower.CL upper.CL
## Mean(others) - Spock    10.7 2.56 39     5.58    15.9
##
## Confidence level used: 0.95
```

Seaweed grazers

```
head(case1301)
```


```
##   Cover Block Treat
## 1    14    B1     C
## 2    23    B1     C
## 3    22    B2     C
## 4    35    B2     C
## 5    67    B3     C
## 6    82    B3     C
```

```
summary(case1301)
```

```
##      Cover      Block      Treat
## Min.   : 1.00   B1      :12   C :16
## 1st Qu.: 9.00   B2      :12   L :16
## Median :22.50   B3      :12   Lf:16
## Mean   :28.62   B4      :12   LfF:16
## 3rd Qu.:42.25   B5      :12   f :16
## Max.   :95.00   B6      :12   fF:16
##                                     (Other):24
```

Seaweed grazers

```
ggplot(case1301, aes(x = Block, y = Cover, shape=Treat, color=Treat)) + geom_jitter(width=0.1, height=0)
```



Seaweed grazers

```

m <- lm(Cover ~ Block + Treat, data = case1301)
summary(m)

##
## Call:
## lm(formula = Cover ~ Block + Treat, data = case1301)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -27.375  -5.812   0.625   5.438  26.125
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)    32.875     4.217   7.795 1.66e-11 ***
## BlockB2         3.750     4.679   0.801 0.425132
## BlockB3        31.750     4.679   6.786 1.59e-09 ***
## BlockB4        45.500     4.679   9.725 2.32e-15 ***
## BlockB5        14.750     4.679   3.153 0.002253 **
## BlockB6        27.750     4.679   5.931 6.67e-08 ***
## BlockB7        13.500     4.679   2.885 0.004980 **
## BlockB8        16.000     4.679   3.420 0.000974 ***
## TreatL        -32.750     4.052  -8.083 4.45e-12 ***
## TreatLf       -35.500     4.052  -8.761 1.96e-13 ***
## TreatLfF     -44.250     4.052 -10.921 < 2e-16 ***
## Treatf        -9.250     4.052  -2.283 0.024995 *
## TreatfF     -18.500     4.052  -4.566 1.71e-05 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##

```

Seaweed grazers

```
anova(m)

## Analysis of Variance Table
##
## Response: Cover
##           Df Sum Sq Mean Sq F value    Pr(>F)
## Block      7  19106   2729.4   20.780 6.977e-16 ***
## Treat      5  23046   4609.1   35.092 < 2.2e-16 ***
## Residuals 83  10902    131.3
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Two-way ANOVA

Regression model with two categorical variable:

$$Y_i \stackrel{ind}{\sim} N(\beta_0 + \underbrace{\beta_1 X_{i,1} + \cdots + \beta_p X_{i,p}}_{\text{variable 1}} + \underbrace{\beta_{p+1} X_{i,p+1} + \cdots + \beta_{p+q} X_{i,p+q}}_{\text{variable 2}}, \sigma^2)$$

where

- $\beta_r, 1 \leq r \leq p$ is the difference between mean response in the reference level compared to the level associated with the r th level
- $\beta_r, p+1 \leq r \leq p+q$ is the difference between mean response in the reference level compared to the level associated with the r th level

F-tests:

1. Variable 1

- Reduced model: no categorical variables $\beta_1 = \cdots = \beta_{p+q} = 0$
- Full model: with first variable $\beta_{p+1} = \cdots = \beta_{p+q} = 0$

2. Variable 2

- Reduced model: with first variable $\beta_{p+1} = \cdots = \beta_{p+q} = 0$
- Full model: with both variables (see model above)

Seaweed grazers

```
anova(m)

## Analysis of Variance Table
##
## Response: Cover
##           Df Sum Sq Mean Sq F value    Pr(>F)
## Block      7  19106   2729.4   20.780 6.977e-16 ***
## Treat      5  23046   4609.1   35.092 < 2.2e-16 ***
## Residuals 83  10902    131.3
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Manuscript statements:

- There is evidence of a difference in mean cover amongst the blocks ($F_{7,83} = 21, p \approx 0$).
- There is evidence of a difference in mean cover amongst the treatments after controlling for blocks ($F_{5,83} = 35, p \approx 0$).

Pygmalion effect

```
head(case1302)
```


```
##   Company      Treat Score
## 1      C1 Pygmalion  80.0
## 2      C1   Control  63.2
## 3      C1   Control  69.2
## 4      C2 Pygmalion  83.9
## 5      C2   Control  63.1
## 6      C2   Control  81.5
```

```
summary(case1302)
```

```
##      Company      Treat      Score
## C1      : 3   Control :19   Min.    :59.50
## C10     : 3   Pygmalion:10  1st Qu.:69.20
## C2      : 3                      Median  :73.90
## C4      : 3                      Mean    :74.07
## C5      : 3                      3rd Qu.:78.90
## C6      : 3                      Max.    :89.80
## (Other):11
```

Pygmalion effect

```
ggplot(case1302, aes(x = Company, y = Score, shape=Treat, color=Treat)) + geom_jitter(width=0.1, height=0)
```



Pygmalion effect

```
m <- lm(Score ~ Company + Treat, data = case1302)
summary(m)

##
## Call:
## lm(formula = Score ~ Company + Treat, data = case1302)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -10.660  -4.147   1.853   3.853   7.740
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   68.39316   3.89308  17.568 8.92e-13 ***
## CompanyC10     4.23333   5.36968   0.788  0.4407
## CompanyC2      5.36667   5.36968   0.999  0.3308
## CompanyC3      0.19658   6.01886   0.033  0.9743
## CompanyC4     -0.96667   5.36968  -0.180  0.8591
## CompanyC5      9.26667   5.36968   1.726  0.1015
## CompanyC6     13.66667   5.36968   2.545  0.0203 *
## CompanyC7     -2.03333   5.36968  -0.379  0.7094
## CompanyC8      0.03333   5.36968   0.006  0.9951
## CompanyC9      1.10000   5.36968   0.205  0.8400
## TreatPygmalion 7.22051   2.57951   2.799  0.0119 *
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 6.576 on 18 degrees of freedom
## Multiple R-squared:  0.5647, Adjusted R-squared:  0.3228
```

Pygmalion effect

```
anova(m)

## Analysis of Variance Table
##
## Response: Score
##           Df Sum Sq Mean Sq F value Pr(>F)
## Company    9  670.98    74.55   1.7238 0.15556
## Treat      1   338.88   338.88   7.8354 0.01186 *
## Residuals 18   778.50    43.25
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Pygmalion Effect

```
anova(m)

## Analysis of Variance Table
##
## Response: Score
##           Df Sum Sq Mean Sq F value Pr(>F)
## Company    9  670.98    74.55  1.7238 0.15556
## Treat      1  338.88   338.88  7.8354 0.01186 *
## Residuals 18  778.50    43.25
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Manuscript statements:

- There is no evidence of a difference in mean score amongst the companies ($F_{9,18} = 2, p = 0.16$).
- There is evidence of a difference in mean score amongst the treatments after controlling for company ($F_{1,18} = 8, p = 0.01$).

Pygmalion Effect

```
drop1(m, test="F")

## Single term deletions
##
## Model:
## Score ~ Company + Treat
##           Df Sum of Sq    RSS    AIC F value    Pr(>F)
## <none>                778.5  117.41
## Company   9     682.52 1461.0  117.67   1.7534  0.14844
## Treat     1     338.88 1117.4  125.89   7.8354  0.01186 *
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Manuscript statements:

- There is no evidence of a difference in mean score amongst the companies **after controlling for treatment** ($F_{9,18} = 2, p = 0.15$).
- There is evidence of a difference in mean score amongst the treatments after controlling for company ($F_{1,18} = 8, p = 0.01$).

ANOVA Tables

- Sequential comparisons
 - Adds new variable to model that already includes variables above it
 - Use `anova()` in R
 - SAS Type I sums of squares
- Partial comparisons
 - Removes variable from model that includes all other variables
 - Use `drop1()` in R
 - SAS Type III sums of squares
 - SPSS default
- Suggestions
 - Results are the same for complete, balanced experiments
 - Always include variables that were part of the experimental design
 - Generally prefer `drop1()`

Three-way ANOVA

```
m <- lm(log(Forrest) ~ Stress + factor(SO2) + factor(O3), data = case1402)
drop1(m, test="F")

## Single term deletions
##
## Model:
## log(Forrest) ~ Stress + factor(SO2) + factor(O3)
##
```

	Df	Sum of Sq	RSS	AIC	F value	Pr(>F)
<none>			0.51159	-106.143		
Stress	1	0.00804	0.51963	-107.675	0.3456	0.5625887
factor(SO2)	2	0.06346	0.57505	-106.635	1.3646	0.2762792
factor(O3)	4	0.75818	1.26977	-86.871	8.1510	0.0003437 ***

```
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Interactions

```
m <- lm(log(Forrest) ~ .^2, data = case1402 %>% select(-William))
drop1(m, test="F")
```

```
## Single term deletions
```

```
##
## Model:
## log(Forrest) ~ (Stress + S02 + 03)^2
##           Df Sum of Sq      RSS      AIC F value Pr(>F)
## <none>                0.50479 -108.54
## Stress:S02  1  0.011613  0.51640 -109.86  0.5291 0.4743
## Stress:03   1  0.012599  0.51739 -109.81  0.5740 0.4564
## S02:03      1  0.014743  0.51953 -109.68  0.6717 0.4209
```