

M6S3 - Pvalue Interpretation

Professor Jarad Niemi

STAT 226 - Iowa State University

October 30, 2018

Outline

- Pvalues
 - Review of calculation procedure
 - Interpretation
- Hypothesis test
 - Decision making
 - Using p-values to make a decision
 - Errors
- ASA Statement on P-values

P-values for $H_0 : \mu = m_0$

Definition

A **p-value** is the (frequency) probability of obtaining a test statistic as or more extreme than you observed if the null hypothesis (model) is true.


So for the null hypothesis $H_0 : \mu = m_0$, calculate

$$t = \frac{\bar{x} - m_0}{s/\sqrt{n}}$$

and find the appropriate probability:

- $H_a : \mu < m_0$ implies $p\text{-value} = P(T_{n-1} < t)$,
- $H_a : \mu > m_0$ implies $p\text{-value} = P(T_{n-1} > t)$, and
- $H_a : \mu \neq m_0$ implies $p\text{-value} = 2P(T_{n-1} > |t|)$,

JMP Example



Alternative Hypothesis	p -value
$H_a : \mu \neq m_0$	0.1677
$H_a : \mu > m_0$	0.9162
$H_a : \mu < m_0$	0.0838

Interpretation

Definition

A ***p*-value** is the probability of obtaining a test statistic as or more extreme than you observed if the null hypothesis is true.

If we assume $X_i \stackrel{iid}{\sim} N(m_0, \sigma^2)$ (because $H_0 : \mu = m_0$) and we have an observed test statistic t based on n observations, the *p*-value is

$$\begin{aligned} P(T_{n-1} < t) & \quad \text{if } H_a : \mu < m_0 \\ P(T_{n-1} > t) & \quad \text{if } H_a : \mu > m_0 \\ P(T_{n-1} > |t| \text{ or } T_{n-1} < -|t|) & \quad \text{if } H_a : \mu \neq m_0 \end{aligned}$$

where

$$T_{n-1} = \frac{\bar{X} - m_0}{S/\sqrt{n}} \sim t_{n-1}$$

which is random because we are considering taking different **random samples** of size n . Thus, the *p*-value is a measure of how often you would expect to see a statistic this extreme if the null hypothesis is true.

Interpretation example

Assume $X_i \stackrel{iid}{\sim} N(\mu, \sigma^2)$ and $H_0 : \mu = 40$. From a random sample with 30 observations you find $\bar{x} = 45.2$ and $s = 11.57$ which results in


$$t = \frac{45.2 - 40}{11.57/\sqrt{30}} \approx 2.462.$$

You have the following probabilities

$$\begin{aligned} P(T_{n-1} > 2.462) &= 0.01 \\ P(T_{n-1} < 2.462) &= 1 - P(T_{n-1} > 2.462) \\ &= 1 - 0.01 \\ &= 0.99 \\ P(T_{n-1} > 2.462 \text{ or } T_{n-1} < -2.462) &= P(T_{n-1} > 2.462) + P(T_{n-1} < -2.462) \\ &= 2P(T_{n-1} > 2.462) \\ &= 1 - 0.01 \\ &= 0.02 \end{aligned}$$

These probabilities correspond to the p -values for the alternative hypotheses $H_a : \mu > 30$, $H_a : \mu < 30$, and $H_a : \mu \neq 30$, respectively.

JMP Example



$$P(T_{35} > -1.4089) = 0.9162$$


$$P(T_{35} < -1.4089) = 0.0838$$

$$P(T_{35} > 1.4089 \text{ or } T_{35} < -1.4089) = 0.1677$$

Hypothesis test for a population mean μ

1. Specify the null and alternative hypothesis.
 - $H_0 : \mu = m_0$ is the default or current belief
 - $H_a : \mu > m_0$ or $\mu < m_0$ or $\mu \neq m_0$
2. Specify a significance level α .
3. Calculate the t -statistic.
4. Calculate the p -value.
5. Make a conclusion:
 - If $p\text{-value} < \alpha$, **reject null hypothesis.**
 - If $p\text{-value} \geq \alpha$, **fail to reject null hypothesis.**


JMP Example



Conclusion at significance level $\alpha = 0.05$:

Alternative Hypothesis	p -value	Conclusion
$H_a : \mu \neq m_0$	0.0522	Fail to reject null hypothesis
$H_a : \mu > m_0$	0.9739	Fail to reject null hypothesis
$H_a : \mu < m_0$	0.0261	Reject null hypothesis

JMP Example



Conclusion at significance level $\alpha = 0.1$:

Alternative Hypothesis	p -value	Conclusion
$H_a : \mu \neq m_0$	0.0522	Reject null hypothesis
$H_a : \mu > m_0$	0.9739	Fail to reject null hypothesis
$H_a : \mu < m_0$	0.0261	Reject null hypothesis

Errors

When performing a hypothesis test, these are the possible situations:

Decision	Truth	
	H_0 true	H_0 not true
Reject H_0	Type I error	correct
Fail to reject H_0	correct	Type II error

Errors:


- A **Type I error** is rejecting the null hypothesis when it is true.
- A **Type II error** is failing to reject the null hypothesis when it is not true.

The **significance level α** is the probability of a Type I error.

ACT scores example

The mean composite score on the ACT among the students at Iowa State University is 24. We wish to know whether the average composite ACT score for business majors is different from the average for the University. Perform a hypothesis test at significance level $\alpha = 0.01$.

Let μ be the average mean composite score amount business majors at Iowa State University. Test $H_0 : \mu = 24$ versus $H_a : \mu \neq 24$.



Reject null hypothesis since $0.002 < 0.01$, i.e. p -value is less than significance level.

ASA Statement on p -values

In 2016, the American Statistical Association published Statement on p -values that states the following principles:

1. P -values can indicate how incompatible the data are with a specified statistical model.
2. P -values do **not** measure the probability that the studied hypothesis is true, or the probability that the data were produced by random chance alone.
3. Scientific conclusions and business or policy decisions should **not** be based only on whether a p -value passes a specific threshold.
4. Proper inference requires full reporting and transparency.
5. A p -value, or statistical significance, does **not** measure the size of an effect or the importance of a result.
6. By itself, a p -value does **not** provide a good measure of evidence regarding a model or hypothesis.

Interpretation (cont.)

The null hypothesis model is

$$X_i \stackrel{iid}{\sim} N(m_0, \sigma^2)$$

for some specified value m_0 .

P -values can indicate how incompatible the data are with [the null hypothesis] model.

The smaller the p -value the larger the incompatibility of the data with the null hypothesis model.

Thus, a small p -value indicates the null hypothesis model is likely not correct. But there are many assumptions in this model that may be wrong, e.g.

- independence,
- identically distributed,
- normality,
- mean is m_0 , and
- constant variance.

The p -value doesn't tell us which one is wrong.